



Bacteriological investigations of skin ulcers in a Swedish pig herd

Karlsson, Frida; Lindberg, Maria; Jensen, Tim Kåre; Pringle, Märit; Fellström, Claes

Publication date:
2014

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):

Karlsson, F., Lindberg, M., Jensen, T. K., Pringle, M., & Fellström, C. (2014). *Bacteriological investigations of skin ulcers in a Swedish pig herd*. Poster session presented at 6th European Symposium of Porcine Health Management, Sorrento, Italy.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Bacteriological investigations of skin ulcers in a Swedish pig herd


Frida Karlsson¹, Maria Lindberg², Tim K. Jensen³, Märit Pringle⁴, Claes Fellström¹

¹Department of Clinical Sciences, Swedish University of Agricultural Sciences, Uppsala, Sweden

²Swedish Animal Health Service, Uppsala, Sweden

³National Veterinary Institute, Technical University of Denmark, Copenhagen, Denmark

⁴National Veterinary Institute, Uppsala, Sweden



Figures a and c showing typical examples of sampled skin ulcers. Figure b showing Fluorescence *In Situ* Hybridization for *Treponema pedis* (red) (CY3) and domain bacteria (green) (FITC) in a section from a porcine skin ulcer F1223. Autofluorescence from erythrocytes is green. Bar 20 μ m.

Conclusions

- Both classic skin pathogens as *S. aureus* and streptococci as well as *Treponema* spp. occur simultaneously in skin ulcers of pigs.
- All isolates of staphylococci and streptococci were susceptible to beta-lactam antibiotics, which are also active against *Treponema* spp.
- Although treponemes were located deep in the skin it is not known whether their role is to lead the way for other skin pathogens or if these bacteria are secondary invaders.

Minimal inhibitory concentration (MIC) mg/L									
of two <i>Staphylococcus aureus</i> isolates and three isolates of β -hemolytic streptococci ^a									
	≤ 0.12	0.25	0.5	1	2	4	8	16	32
Penicillin	S.a. ^b S. β . ^c								
Ampicillin				S.a. S. β .					
Ceftiofur		S. β .		S.a.					
Spiramycin						S.a. S. β .			
Neomycin						S.a.			(S. β .) ^d
Gentamicin				S.a.		(S. β .)			
Streptomycin						S.a.1 ^e		S.a.3 ^f	(S. β .)
Trim./Sulph. ^g			S.a. S. β .						
Enrofloxacin		S.a.		(S. β .)					
Tetracycline			S.a.		S. β .7 ^h			S. β .3,4 ⁱ	
Florfenicol				S. β .		S.a.			
Oxacillin			S.a. S. β .						

^a The white fields denote range of dilutions tested for each substance. MICs above the range are given as the concentration closest to the range. MICs equal to or lower than the lowest concentration tested are given as the lowest tested concentration. Bold vertical lines indicate cut-off values defining resistance according to VetMIC Large Animal, National Veterinary Institute, Sweden. ^b S.a.: *Staphylococcus aureus* isolates 1221 and 1223. ^c S. β : β -hemolytic streptococci isolates 1223, 1224 and 1227. ^d (S. β): Not relevant as the inherent susceptibility is above concentrations that can be obtained during therapy. ^e S.a.1: *Staphylococcus aureus* isolate 1221. ^f S.a.3: *Staphylococcus aureus* isolate 1223. ^g Concentration of trimethoprim given, tested in concentration ratio 1/20 (trimethoprim/sulphamethoxazole). ^h S. β .7: β -hemolytic streptococci isolate 1227. ⁱ S. β .3, 4: β -hemolytic streptococci isolates 1223 and 1224.

Overview of samples and methods					
Sample	W-S ^a	FISH		PCR ^b (<i>T. spp</i>)	Aerob culture
		<i>T. spp</i>	<i>T. pedis</i>		
E ^c 1221	+	+	+	+	Nt ^d
E 1222	-	-	Nt	-	Nt
E 1223	-	+	+	-	Nt
E 1224	-	+	+	+	Nt
E 1225	-	-	Nt	-	Nt
F ^e 1221	-	-	Nt	Nt	S.a. ^f , S. β . ^g
F 1223	-	+	+	Nt	S.a., S. β .
F 1224	-	+	+	Nt	S.a., S. β .
F 1226	Nt	Nt	Nt	-	S.a., S. β .
F 1227	Nt	Nt	Nt	-	S.a., S. β .

W-S^a: Warthin Starry silver staining, PCR^b: Intergenic spacer region 2- based PCR, E^c: Ear necrosis, Nt^d: not tested, F^e: other type of skin ulcer, S.a.^f: *Staphylococcus aureus*, S. β .^g: β -hemolytic streptococci

Aims

- Investigate the presence of *S. aureus*, streptococci and *Treponema* spp. in skin ulcers of pigs in one herd.
- Test the antimicrobial susceptibility of obtained isolates of *S. aureus* and streptococci.

Acknowledgement

This project was supported by Formas (Swedish Research Council for Environment, Agricultural Sciences and Spatial Planning).